

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Synthetic cathinone adulteration of illegal drugs.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1730609> since 2020-02-25T11:37:28Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Psychopharmacology

Synthetic Cathinone Adulteration of Illegal Drugs

--Manuscript Draft--

Manuscript Number:	PSPH-D-18-00256R1	
Full Title:	Synthetic Cathinone Adulteration of Illegal Drugs	
Article Type:	Review	
Funding Information:	National Institute on Drug Abuse (R01 DA039139)	Dr. Scott Rawls
Abstract:	<p>Abstract</p> <p>Rationale Current prevalence estimates of synthetic cathinone ("bath salts") use may be underestimates given that traditional metrics (e.g., surveys, urinalysis) often fail to capture the emergent issue of synthetic cathinone adulteration of more common illegal drugs such as ecstasy.</p> <p>Objectives This review examines the evolution of synthetic cathinones and prevalence of use over the past decade in the United States. We also review methods of self-report and biological testing of these compounds, and adverse outcomes pertaining to adulterated drug use.</p> <p>Results Synthetic cathinone use emerged in the US by 2010 with use associated with tens of thousands of poisonings. Reported poisonings and self-reported use have substantially decreased over the past five years. However, our review suggests that current estimates of use are underestimates due to underreporting, stemming primarily from unknown or unintentional use of adulterated formulations of illegal drugs such as ecstasy. Unintentional synthetic cathinone use can also be associated with adverse outcomes reportedly from use of other drugs individuals believed they were using.</p> <p>Conclusions While intentional synthetic cathinone use has decreased in recent years, evidence suggests prevalence of use is underestimated. Testing of drugs and/or biological specimens can help us better determine exposure to synthetic cathinones and researchers and clinicians should become better aware that unintentional exposure to these potent compounds (e.g., as adulterants) is often unknown or unintentional. Research utilizing both survey and biological testing methods may help more accurately determine the prevalence of use of these compounds.</p>	
Corresponding Author:	Chicora Frances Oliver, B.A. Temple University Philadelphia, PA UNITED STATES	
Corresponding Author Secondary Information:		
Corresponding Author's Institution:	Temple University	
Corresponding Author's Secondary Institution:		
First Author:	Chicora Frances Oliver, B.A.	
First Author Secondary Information:		
Order of Authors Secondary Information:		
Order of Authors:	Chicora Frances Oliver, B.A. Joseph Palamar, PhD Alberto Salomone, PhD Steven James Simmons, PhD Helene Philogene-Khalid, PhD Nick Stokes McCloskey, B.S. Scott Rawls, PhD	
Author Comments:		

Response Letter.

Re: Reviewer response for manuscript
Klaus A. Miczek PhD
Principal Editor of *Psychopharmacology*

Dear Dr. Miczek --

Sincere thanks to you and the reviewing team for considering our manuscript for publication in *Psychopharmacology*. We have addressed each comment, and an itemized reply can be found below this passage. The reviewer comment will be **boldfaced**, and our reply will follow.

We are happy to make additional edits to the manuscript as you and the reviewing team see fit.

Sincerely,
Chicora Oliver.

Editorial Comment

We have now included a Table that specifies effects of synthetic cathinones administered with (or following) other illicit drugs including cocaine and methamphetamine. We believe this Table is a valuable reference that will refer readers to original research reports addressing this critical issue. We regretfully were unable to construct a timeline of historical development of synthetic cathinones but hope the reviewing team finds the written text to be sufficient.

Reviewer 1

(1-2) **“The review comes to a little bit of an abrupt end... (add) recommendation for professional education (ED personnel)...”**. We thank Reviewer 1 for this comment and have expanded our concluding passages. Specifically, we reiterate take-home messages that drugs marketed as something other than synthetic cathinones may, indeed, contain one or more synthetic cathinone agents. We expand on potential factors contributing to suicides associated with synthetic cathinone use.

All minor corrections were performed in the revised manuscript.

Reviewer 2

(1-2) **“In the abstract...identify "ecstasy" by its chemical name.” and “Page 9, line 10 - please define "ersatz".”** We thank Reviewer 2 for noting these issues with clarity. We define ecstasy in the abstract and use the clearer term “counterfeit” in place of “ersatz”.

(3) **“Toxicological Assessment subsection - it should be mentioned that references standards for many parent compounds are now available (i.e., Cayman Chemical, Cerillian). Also, in this section it should be mentioned that GC-MS methods are also used for detection and analysis.”** These important changes were made to the revised text. Discussion of toxicological assessment now cites the availability of reference standards as well as GC-MS testing methods.

(4) **“ It would be helpful to know, if such information is available, the degree to which some drugs such as MDMA are adulterated with cathinones. For example, in seized MDMA tablets, what percent of the tablets are MDMA vs. cathinones? Likely there are trace amounts, but the relative contamination levels could be useful to discuss briefly.”** We agree with Reviewer 2 that having this information would be incredibly useful. However, such data is not currently available. The coauthors have personally contacted likely sources of this information (e.g. NFLIS) but have been unable to find data relevant to this issue.

(5) **“Consequences subsection - when discussing the ability of cathinones to increase, DA, NE, 5-HT, the authors should briefly clarify that these increases are in extracellular levels, not tissue content, and are achieved by presynaptic transporter blockade or via acting as transporter substrates. The phrase "increase dopamine, norepinephrine, and serotonin..." is too general.”** This important oversight is corrected in the revised text. We made

clear that neurotransmitters modulated by synthetic cathinones are done so in the synapse and result from both transporter blocking and substrate properties of these drugs.

(6) **“The final Conclusions section is very short and could be expanded with a discussion of some critical issues. These could include, but are not limited to, the issue of constant development of new cathinone derivatives, identification of synthetic cathinone metabolites which might have long elimination half lives which better aid in detection in biological samples as markers of prior use, identification of unique behavioral or physiological biomarkers of contaminated drug use, etc.”** We thank Reviewer 2 for this incredibly helpful feedback. The Conclusion (now “Concluding Remarks”) was expanded to discuss the potential for death from overdose or fatal self-harm and how there is a need for more awareness of the symptoms of synthetic cathinone overdose. We also discuss the need for more research on the behavioral and neurotoxic effects of synthetic cathinone drug combinations. We also highlight the need for studies that examine how environmental factors such as temperature alter the effects of synthetic cathinones use. Finally, we close by stating that these efforts may aide in the development of interventions that address the unique problem of synthetic cathinone adulteration of illegal drugs. We believe these changes greatly strengthen the text and hope the reviewing team find these changes to be sufficient.

[Click here to view linked References](#)

Synthetic Cathinone Adulteration of Illegal Drugs

Chicora F. Oliver^{*1}, Joseph J. Palamar², Alberto Salomone³, Steven J. Simmons^{1,5}, Helene Philogene-Khalid⁴, Nick Stokes-McCloskey¹, Scott M. Rawls¹

¹Center for Substance Abuse Research, Lewis Katz School of Medicine at Temple University, 3500 North Broad Street, 882A, Philadelphia, PA 19140, USA.

²Department of Population Health, 227 E. 30th Street, 7th Floor, New York, NY 10016, USA.

³Centro Regionale Antidoping "A. Bertinaria", Regione Gonzole 10/1, 10043 Orbassano, Torino, Italy.

⁴Department of Psychiatry, 3400 Spruce Street, Philadelphia, PA 19104.

⁵Department of Anesthesiology and Critical Care Medicine, Children's Hospital of Philadelphia Research Institute, Philadelphia, PA, 19104.

Conflicts of Interest: There are no conflicts of interest to report.

Acknowledgement of funding: National Institutes of Health, National Institute on Drug Abuse R01 DA039139 to SMR. SJS was supported by institutional training fellowships (T32 DA007237 to Ellen M. Unterwald, T32 NS007413 to Michael B. Robinson).

***Corresponding Author**

Chicora F. Oliver

Center for Substance Abuse Research

Lewis Katz School of Medicine at Temple University

Email: chicoraoliver@temple.edu

Phone: 215-707-3656

Fax: 215-707-6932

1
2
3
4 **Abstract.**
5

6 *Rationale.* Current prevalence estimates of synthetic cathinone (“bath salts”) use may be
7
8 underestimates given that traditional metrics (e.g., surveys, urinalysis) often fail to capture the
9
10 emergent issue of synthetic cathinone adulteration of more common illegal drugs such as ecstasy
11
12 (3,4-methylenedioxymethamphetamine).
13
14

15
16 *Objectives.* This review examines the evolution of synthetic cathinones and prevalence of use
17
18 over the past decade in the United States. We also review methods of self-report and biological
19
20 testing of these compounds as well as adverse outcomes associated with adulterated drug use.
21
22

23
24 *Results.* Synthetic cathinone use emerged in the United States by 2009 with use associated with
25
26 tens of thousands of poisonings. Reported poisonings and self-reported use have substantially
27
28 decreased over the past five years. However, our review suggests that current estimates of use
29
30 are underestimates due to underreporting stemming primarily from unknown or unintentional use
31
32 of adulterated formulations of relatively popular illegal drugs, such as ecstasy.
33
34

35
36 *Conclusions.* While intentional synthetic cathinone use has decreased in recent years, evidence
37
38 suggests prevalence of use is underestimated. Testing of drugs and/or biological specimens can
39
40 improve the accuracy of synthetic cathinone use estimates. Furthermore, we advocate that
41
42 researchers and clinicians should become better aware that exposure to these potent compounds
43
44 (e.g., as adulterants) often occurs unknowingly or unintentionally. To improve our understanding
45
46 of synthetic cathinone adulteration, research utilizing a combinatorial approach (survey and
47
48 biological testing) will help more accurately determine the prevalence and impact of this public
49
50 health issue.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 **Keywords.**
5

6
7 Synthetic cathinones; novel psychoactive substances; ecstasy; adulteration; addiction.
8
9

10
11 **Abbreviations.**
12

13
14 α -PVP - α -pyrrolidinopentiophenone
15

16
17 4-MMC - 4-methylmethcathinone
18

19
20 5-HT - serotonin
21

22
23 AAPCC - American Association of Poison Control Center
24

25
26 DA - dopamine
27

28
29 DEA - Drug Enforcement Administration
30

31
32 ED – emergency department
33

34
35 EDM - electronic dance music
36

37
38 MDMA - 3,4-methylenedioxymethamphetamine
39

40
41 MDMC - 3,4-methylenedioxymethcathinone
42

43
44 MDPV – 3,4-methylenedioxypyrovalerone
45

46
47 NA - noradrenaline
48

49
50 NFLIS - The National Forensic Laboratory Information System
51

52
53 NPS – new psychoactive substance
54

55
56 PCP – phencyclidine
57

58
59 UHPLC-HRMS - ultra high performance-tandem mass spectrometry-high resolution mass
60
61 spectrometry
62

63
64 UHPLC-MS/MS - ultra high performance liquid chromatography-tandem mass spectrometry
65

Synthetic Cathinone Adulteration of Illegal Drugs.

A growing number of studies have demonstrated synthetic cathinone adulteration of illegal drugs such as ecstasy or 3,4-methylenedioxymethamphetamine (MDMA, “ecstasy”) (Rust et al. 2012; Palamar et al. 2016d, 2017b; Salomone et al. 2017). Various synthetic cathinones (colloquially, “bath salts”) produce effects similar to those of MDMA, and therefore adulteration or replacement of ecstasy with synthetic cathinones may go unnoticed by some users. However, synthetic cathinones can be much more potent than the drugs they are intended to mimic (Baumann et al. 2012), and thus the risk of overdose and death can be greatly increased. While growing awareness, legal restrictions, and sensational media reporting seem to have reduced intentional synthetic cathinone use, synthetic cathinones continue to be detected in users unaware of consumption. For example, two recent studies analyzing hair samples of ecstasy-using electronic dance music (EDM) party attendees found that 50-68% of those testing positive for synthetic cathinones denied known use (Palamar et al. 2016c, 2017c). Retrospective analysis of the hair of drug users confirms the presence of synthetic cathinones in those who report MDMA and/or other drug use but deny conscious consumption of synthetic cathinones (Rust et al. 2012; Palamar et al. 2016c, 2017c). These and other studies reviewed in this manuscript indicate that current prevalence estimates of synthetic cathinone use (e.g., in the United States) are likely underestimates, and we believe addressing these potential underestimates is critical for improving public health and safety.

Evolution of Synthetic Cathinone Use.

1
2
3
4 Synthetic cathinones are psychoactive substances manufactured to mimic the stimulating
5
6 effects of naturally occurring, plant-based cathinones. The psychoactive properties of cathinones
7
8 have been appreciated for several millennia through the practice of chewing leaves of the
9
10 cultivated perennial shrub khat, *C. edulis* (native to East Africa/Arabian Peninsula). Subjective
11
12 effects of khat leaf chewing include allayed fatigue and elated mood (Alles et al. 1961; Numan
13
14 2004). In the mid-20th century, means of expedited international trade allowed perishable khat to
15
16 be exported to the United States, United Kingdom and elsewhere. The United Nations Narcotics
17
18 Laboratory successfully isolated (\pm)-cathinone as the principal psychoactive agent from khat
19
20 leaves in the early 1970s (for discussion, see Kalix 1984) and noted structural similarity to
21
22 phenylisopropylamine (amphetamine) with the addition of a ketone at cathinone's β -carbon
23
24 atom. In 1987, Glennon and colleagues (1987) discovered alarming potency in the amphetamine-
25
26 like subjective effects of methcathinone among animals, although readers should note the
27
28 original synthesis of methcathinone was performed in the 1920s in an effort to produce ephedrine
29
30 homologs (Hyde et al. 1928). In 1994, the Drug Enforcement Agency (DEA) of the United States
31
32 designated methcathinone as a Schedule I illegal drug (State Department 1994), and use of
33
34 synthesized cathinones remained low for the subsequent two decades.
35
36
37
38
39
40
41
42

43 Cathinone-derived psychostimulant drugs surged in popularity in the late 2000s most
44
45 notably across the United Kingdom and United States. Novel modifications of the parent
46
47 compound yielded abuse-prone agents including amphetamine-like 4-methylmethcathinone (4-
48
49 MMC, "mephedrone") and 3,4-methylenedioxypyrovalerone (MDPV), as well as the
50
51 empathogenic agent 3,4-methylenedioxymethcathinone (MDMC, "methylone"), which
52
53 structurally resembles MDMA. The surge in use is attributed to marketing on the "dark web" as
54
55 novel formulations obviated legal persecution for several years by classifying agents as "not for
56
57
58
59
60
61
62
63
64
65

human consumption” with product names including “blue silk,” “vanilla sky,” and “white lightning” (European Monitoring Centre for Drugs and Drug Addiction 2015; Drug Enforcement Agency 2017). The National Forensic Laboratory Information System (NFLIS) reported 34 analytically-confirmed cases of synthetic cathinone seizures across 8 states in 2009. In 2010, NFLIS reported 628 cases across 27 states (NFLIS 2011), and by July 2012, mephedrone and MDPV had become classified as Schedule I by the DEA under emergency order. Consequently, NFLIS reported methylone as the most prevalent synthetic cathinone in 2013 (12,067 cases, 71.8% of all synthetic cathinone cases; NFLIS 2015).

“Second generation” synthetic cathinones surfaced after methylone was scheduled in April 2013 and included 3,4-methylenedioxy-*N*-ethylcathinone (ethylone) and α -pyrrolidinopentiophenone (α -PVP). Synthetic cathinones seized in this “second generation” era possessed ethyl, butyl, pyrrolidinyl, fluoro and benzyl substitutions against cathinone’s backbone (for review, see Majchrzak et al. 2018). Use of the aforementioned synthetic cathinones is associated with health risks ranging from injection site irritation/infection to paranoia/hallucinatory delirium, sympathomimetic toxicity, serotonin syndrome, and sudden cardiac death (Wood et al. 2010; Dorairaj et al. 2012; Carbone et al. 2013). α -PVP, commonly referred to as “flakka,” was linked to 63 deaths in Fort Lauderdale, Florida within a 16-month period (CBS 2016), and the bizarre behavior linked with its use has earned it the street name “zombie drug” (Nóbrega and Dinis-Oliveira 2018). Data as of mid-2017 show β -keto-ethylbenzodioxolypentamine (*N*-ethylpentylone, “ephylone”), which appears to possess appreciably greater potency than MDMA, is the most widely used synthetic cathinone in the United States, with over 2,500 cases reported (DEA 2017; Krotulski et al. 2018). Furthermore, 13 individuals were hospitalized in New Zealand from mistakenly consuming high doses of

1
2
3
4 ephylone in March 2018 (Newshubb 2018). These recent data confirm that novel synthetic
5
6 cathinones (administered alone or in formulation) continue to be produced and thus pose a threat
7
8 to public health on an international scale. And while deliberate abuse of synthetic cathinones has
9
10 ostensibly declined, the variability of drug adulteration obscures the public's real rates of
11
12 consumption.
13
14

15 16 **Epidemiology of Synthetic Cathinone Use.** 17

18
19 The American Association of Poison Control Center's (AAPCC) National Poison Data
20
21 System tracks reported poisonings throughout the United States. AAPCC did not begin receiving
22
23 reports of exposure to synthetic cathinones until 2009 - this first year, AAPCC received 298
24
25 cases of synthetic cathinone poisonings (Wood 2013). In 2011, reports of exposures increased to
26
27 >6,000 (Wood 2013), and synthetic cathinones were involved in over 20,000 drug-related
28
29 emergency department (ED) visits (Substance Abuse and Mental Health Services Administration
30
31 2011). Between 2011 and 2013, there were nearly 10,000 reported exposures to synthetic
32
33 cathinones; however, exposures decreased to fewer than 1,000 between 2014 and mid-2015
34
35 (Wood 2013; AAPCC, 2016). Recent public data are not available on hospitalizations or
36
37 poisonings involving synthetic cathinones in the United States for recent years, but there is
38
39 evidence to suggest diminution. While national drug-related ED data after 2011 are not available,
40
41 it should be noted that in light of the prevalence of ecstasy use decreasing dramatically after the
42
43 early 2000s (Johnston et al. 2018), ED visits related to "ecstasy" use increased several-fold from
44
45 4,452 in 2001 to 22,498 in 2011 (Substance Abuse and Mental Health Services Administration
46
47 2011). The steep increase in "ecstasy"-related ED visits through 2011 coincided with popularity
48
49 of use of the drug in powder form (referred to as "Molly"), which may increase likelihood of
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 adulteration (Palamar 2017). It is unknown to what extent synthetic cathinones were
5
6
7 unknowingly ingested by users as toxicology testing was not utilized in EDs.
8

9 Monitoring the Future (MTF), an annual nationally representative survey of high school
10
11 seniors (12th graders), is the only national survey in the US to query “bath salts” (synthetic
12
13 cathinone) use. MTF first asked about “bath salts” in 2012, and 1.3% of students reported past-
14
15 year use (Johnston et al. 2018). Prevalence of past-year use appears to have decreased with 0.6%
16
17 reporting use in 2017 (Palamar 2015; Johnston et al. 2018). However, a major limitation of the
18
19 MTF survey is that “bath salts” are not clearly defined, nor are examples of specific compounds
20
21 provided. Few studies have examined synthetic cathinone use in greater depth. In 2016, a study
22
23 of EDM party attendees tested survey methods querying synthetic cathinone use and found that
24
25 nearly a tenth (9.3%) of those denying overall use of synthetic cathinones then reported use of a
26
27 specific synthetic cathinone on the following page (Palamar et al. 2017a), reflecting widespread
28
29 confusion as to what drugs constitute “bath salts.” This was most commonly the case with
30
31 methylone and mephedrone, which are synthetic cathinones with empathomimetic, ecstasy-like
32
33 effects. Further, an analysis of type-in responses in the National Survey of Drug Use and Health
34
35 found that type-in responses of other drugs used led to a severe underestimate of only 0.05% of
36
37 individuals in the United States ever using synthetic cathinones or other novel phenethylamines
38
39 or euphorigenic stimulants (Palamar et al. 2015b). These findings reveal that more commonly
40
41 employed means of assessing drug use (e.g., surveys) may underestimate synthetic cathinone
42
43 use, in part, due to lack of drug education among users. Few other studies have queried synthetic
44
45 cathinone use on surveys. One survey of college students in 2011-2012 included a question about
46
47 use of synthetic cathinones and a specific compound (MDPV), and use was reported by only 1%
48
49 of the sample (Miller and Stogner 2014). Similarly, a survey of attendees of gay clubs, hip hop
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 clubs, indie rock clubs, and EDM clubs in New York City (NYC) in 2012 asked about use of
5
6 synthetic cathinones or mephedrone, and similarly, only 1% reported use (Kelly et al. 2013).
7
8

9 Surveys focusing on EDM party attendees, however, have tended to find higher
10
11 prevalence of reported synthetic cathinone use. We believe higher prevalence is detected in this
12
13 population for four reasons. First, EDM party attendees are at high risk for using various drugs
14
15 (Ross et al. 2003; Palamar et al. 2015a, 2017b; Kurtz et al. 2017), and use of other drugs has
16
17 been found to be a major risk factor for synthetic cathinone abuse (Palamar et al. 2015b, 2017b;
18
19 Fernández-Calderón et al. 2018). Second, various synthetic cathinone compounds have similar
20
21 effects to common club drugs such as ecstasy, with some compounds like mephedrone
22
23 functioning as counterfeit replacements (Brunt et al. 2011a). Third, surveys of EDM party
24
25 attendees tend to ask about multiple synthetic cathinone compounds and not just synthetic
26
27 cathinones as an overall category, and it appears that asking about multiple compounds increases
28
29 the likelihood of recognizing a specific compound that was used. Finally, EDM party attendees
30
31 have been found to be at high risk for unknowingly or unintentionally using synthetic cathinones
32
33 as these compounds are common adulterants in drugs such as ecstasy (Palamar et al. 2016c,
34
35 2017c).
36
37
38
39
40
41
42

43 A self-selected online sample of past-year drug-using nightclub attendees who
44
45 participated in the Global Drug Survey in 2013 found that one-out-of-ten (10%) reported ever
46
47 using at least one synthetic cathinone queried (Palamar et al. 2016b). Most (8%) reported
48
49 methylone use, 5% reported mephedrone use, and 2% reported MDPV use. More recently, time-
50
51 space sampling and weighting methods have been used by some studies to yield more accurate
52
53 estimates of synthetic cathinone use from venue-based recruitment in the EDM scene. A study of
54
55 young adult (age 18-25) EDM party attendees in 2015 estimated that 7% of attendees had used at
56
57
58
59
60
61
62
63
64
65

1
2
3
4 least one (of 26) synthetic cathinone compound, with methylone (3%) being the most prevalent
5
6 (Palamar et al. 2016a). In 2016, similar lifetime prevalence of 7.7% was estimated among EDM
7
8 attendees in NYC age 18-40 (Palamar et al. 2018); however, in a new 2017 cohort of attendees,
9
10 estrimated prevalence of known use decreased in half to 3.5% (Palamar 2018).
11
12
13

14 Prevalence estimates of synthetic cathinone use also tend to be underestimates because
15
16 many users of drugs such as ecstasy consume these compounds unknowingly, as synthetic
17
18 cathinones are common adulterants or replacements (discussed in detail below). In fact, during a
19
20 recent survey of EDM party attendees in NYC, 30% of respondents agreed that “bath salts” are
21
22 sometimes found in Molly, and 15% reported believing they might have unknowingly used “bath
23
24 salts” (Palamar 2018). This suggests many EDM attendees are aware of the high potential for
25
26 adulteration, hence extensive interest in “drug checking” in which an individual’s drug is tested
27
28 for purity (Barratt et al. 2018). To our knowledge, no studies have examined prevalence of
29
30 adulteration in the United States, and this is likely due to high legal risk for both users and drug
31
32 checkers. However, studies have begun to analyze biological specimens of users to determine
33
34 whether they have used synthetic cathinones.
35
36
37
38
39
40

41 **Toxicological Assessment of Synthetic Cathinone Use.**

42

43 After the presence of synthetic cathinones on the black market became recognized,
44
45 forensic and clinical laboratories worldwide had to face the analytical challenge of identifying
46
47 and quantifying these new drugs in various biological matrices. The commercial unavailability of
48
49 reference standards for the parent drugs and their metabolites, the lack of updated and
50
51 comprehensive immunoassays for their detection, and the insufficient investigation of their
52
53 metabolic transformation after intake represented just the main problems toward the
54
55 identification and quantification of synthetic cathinones (Salomone et al. 2017). At the moment,
56
57
58
59
60
61
62
63
64
65

1
2
3
4 most of the aforementioned challenges have been at least partially overcome. For example,
5
6 reference standards for many parent compounds are now available from companies such as
7
8 Cayman Chemical and Cerillian. Gas chromatography with mass spectrometry (GC-MS) is also
9
10 employed to assess synthetic cathinone use (Alvarez et al. 2017). The most recent equipment
11
12 based on ultra high performance-tandem mass spectrometry (UHPLC-MS/MS) and high
13
14 resolution mass spectrometry (UHPLC-HRMS) allows the detection of expansive synthetic
15
16 cathinone panels within a single analytical run. Several laboratories currently embrace this
17
18 promising advent by offering screening and confirmation analysis for synthetic cathinones (Zuba
19
20 and Adamowicz 2018; Marchei et al. 2018).

21
22
23
24
25
26 Much is yet to be discovered about the current diffusion of synthetic cathinones among
27
28 the general population and in high-risk populations. Synthetic cathinones are eliminated from the
29
30 urine, blood, and saliva of users within hours or days, which limits the ability of toxicological
31
32 confirmation. This is especially concerning in cases of hospitalization and death resulting from
33
34 intentional or unintentional use of synthetic cathinones (Salomone et al. 2017). As a
35
36 consequence, a number of unresolved issues are raised, including the number and variety of
37
38 synthetic cathinones present in different countries, their frequency of use, and user
39
40 demographics. To circumvent the limitations of conventional biological samples (e.g., blood,
41
42 urine, saliva), the detection of synthetic cathinones in alternative matrices (i.e., hair samples) was
43
44 proposed as a practical means to provide preliminary information on the black market
45
46 penetration of synthetic cathinones (Rust et al. 2012; Salomone et al. 2016b) in specific
47
48 territories and populations. The keratin matrix incorporates parent synthetic cathinones
49
50 consumed over extended time periods, providing access to a much wider diagnostic window than
51
52 blood and urine. This feature, combined with the analytical performance of the latest generation
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 of UHPLC-MS/MS instruments, allows researchers to obtain significant information about past
5
6 use – even a single intake of any targeted compound, with older periods of use corresponding to
7
8 the hair segments more distant from the hair root (Salomone et al. 2016b). In particular,
9
10 analytical methods must guarantee high sensitivity: due to their high pharmacological potency *in*
11
12 *vitro*, it is likely that synthetic cathinones are active in relatively low doses (and therefore are
13
14 consumed in low doses, and present at lower levels in hair).
15
16
17

18
19 Remarkably, the majority of the published studies describing novel synthetic drug
20
21 detection in hair samples reported the frequent occurrence of polydrug use (Kintz et al. 2015;
22
23 Salomone et al. 2016a). One explanation for this is because the consumers are unaware of the
24
25 actual composition of the products. Today, despite the general knowledge that no systematic
26
27 correspondence exists between the trade or the street name and real content of the
28
29 pill/powder/crystal/herbal blend, consumers hardly ever have means of verifying what they
30
31 purchased. This is not surprising as synthetic cathinones are semi-clandestine preparations in
32
33 which the active ingredients may vary over time. This inconsistency in the content of illegal
34
35 drugs implies that consumers do not have control of the potency and efficacy of the products to
36
37 be consumed. Even when synthetic cathinones are purchased online or used intentionally,
38
39 substantial risk exists that they are mislabeled either because they contain chemical analogs of
40
41 the ordered drug (e.g., pentedrone instead of 3,4-dimethylcathinone) or because the active agent
42
43 differs from what was advertised by the dealer or via the website (Brunt et al. 2017). In this
44
45 scenario, hair testing offers a unique perspective in the investigation of drug consumption,
46
47 provided a large panel of target analytes. The extended diagnostic time window covered by the
48
49 keratin matrix (unlike urine, blood, and saliva) allows retrospective investigation of drug
50
51 prevalence and diffusion of any targeted psychoactive substance. In particular, the occurrence of
52
53
54
55
56
57
58
59
60
61
62
63
64
65

poly-use and unintentional intake of synthetic cathinones can be revealed by means of hair analysis, possibly combined with surveys.

Synthetic Cathinone Adulteration of Illegal Drugs.

Rust et al. (2012) examined hair samples from 2009 to 2010 from individuals who originally tested positive for amphetamines and/or MDMA (n=325) (Rust et al. 2012). Hair was collected as part of a driving ability assessment and employed liquid chromatography with tandem mass spectrometry (LC-MS/MS) to test for a variety of novel psychoactive substances (NPS), including synthetic cathinones. Of the 120 samples that tested positive for NPS, 10% tested positive for synthetic cathinones (11 samples for mephedrone, 1 sample for methylone). A later study also found high levels of mephedrone in suspected drug users. In France, Martin et al. (2012) found that nearly 20% of hair samples collected from those suspected of narcotic abuse (13 out of 67) tested positive for mephedrone (Martin et al. 2012). While it is unclear whether individuals in either of these studies were aware of synthetic cathinone use, these methods confirm the utility of analyzing hair to assess synthetic cathinone use in suspected users.

In 2015, Palamar and colleagues surveyed and analyzed hair samples from 48 EDM attendees in NYC reporting lifetime ecstasy/MDMA/Molly use (Palamar et al. 2016c). Almost half (48%) of hair samples contained butylone and 10% contained methylone. However, of those who reported no lifetime use of synthetic cathinones or unknown pills or powders, 41% tested positive for a synthetic cathinone or other NPS. Specifically, of those denying use of any synthetic cathinones listed on the survey, 38% tested positive for butylone, 9% for methylone, and 3% for the highly potent compound α -PVP (“Flakka”). This study also found that racial/ethnic minorities were more likely to test positive for synthetic cathinones after denying lifetime use. The strongest predictor of testing positive for synthetic cathinones was frequent nightclub/festival attendance.

1
2
3
4 In 2016, a similar study was conducted with a focus on 90 past-year ecstasy users in the
5
6 NYC EDM scene (Palamar et al. 2017c). While over a quarter (28%) of attendees tested positive
7
8 for synthetic cathinones, 68% of those testing positive denied past-year use. Butylone (14%),
9
10 ethylone (11%), and pentylone (10%) were the most commonly detected synthetic cathinones -
11
12 strikingly, all participants denied use of these compounds. Methylone was detected in 3% of hair
13
14 samples, but two of these three participants reported known use of this substance. Unknown α -
15
16 PVP use was also detected in 2% of samples. These studies demonstrate that many ecstasy-using
17
18 nightclub/festival attendees are unintentionally administering synthetic cathinones, and that drug
19
20 checking may be beneficial for those unwilling or unable to abstain.
21
22
23
24

25
26 Synthetic cathinone use at EDM parties has also led to clusters of hospitalizations. In
27
28 particular, there were at least 22 drug-related hospitalizations at the 2013 Electric Zoo EDM
29
30 festival in NYC (Ridpath et al. 2014). Of the 22 reported hospitalizations, 9 (41%) were
31
32 classified as severe, and there were two deaths. Of the 22 patients, 17 were administered
33
34 toxicological tests, among whom 55% had used “synthetic club drugs.” Four of the 17 tested
35
36 positive for methylone alone, 3 for methylone and MDMA, 2 for methylone and other illegal
37
38 drugs other than MDMA, and 2 for MDMA alone. One decedent tested positive for MDMA,
39
40 while the other tested positive for both MDMA and methylone. It is unclear whether the second
41
42 decedent or any of the other patients who tested positive for methylone were fully aware of their
43
44 drug/formulation intake.
45
46
47
48
49

50
51 Various drug-testing services throughout Europe have also detected a range of synthetic
52
53 cathinones in drugs sold as ecstasy. For example, despite the increasing purity of ecstasy tested
54
55 from 2008-2013 (Brunt et al. 2017), a range of synthetic cathinones including 4- MEC,
56
57 mephedrone, methylone, and ethylone were detected in submitted substances. The Drug
58
59
60
61
62
63
64
65

1
2
3
4 Information and Monitoring System in the Netherlands in particular detected mephedrone sold as
5
6
7 ecstasy in a large portion of samples throughout 2009 (Brunt et al. 2011b). A Spanish drug-
8
9
10 testing study also examined drug submissions from 2010-2012 and detected 22 different
11
12 synthetic cathinone compounds (primarily methylone, mephedrone, 4- MEC, and MDPV) in
13
14 drugs to be sold as ecstasy or ketamine (Caudevilla-Gállico et al. 2013).

15
16 It should be noted that while advanced laboratory methods are now able to detect the
17
18 presence of many synthetic cathinones, a major limitation to research is that standard drug tests
19
20 (e.g., urine dip sticks) cannot detect the presence of several dangerous substances. Without a
21
22 simple form of testing, it is difficult for both researchers and medical professionals (e.g., ED
23
24 staff) to test body fluids and determine potential exposure to these compounds. Therefore,
25
26 underestimation of use may continue. However, some harm reduction organizations such as
27
28 DanceSafe sell drug testing reagents which provide users a color-change metric of purity.
29
30
31 Reagent tests are not the most reliable form of testing, but they can improve ability to detect drug
32
33 adulteration. On-site (e.g., at dance festivals) or off-site testing (e.g., at analytical facilities) can
34
35 help track what drugs people are really using (Butterfield et al. 2016), and this could identify
36
37 specific synthetic cathinones and help detect trends in exposure. Most importantly, drug testing
38
39 is an opportunity for users to discard the drug they plan to use, and results can inform future use
40
41 or decisions to purchase from specific individuals (Butterfield et al. 2016).
42
43
44
45
46
47

48 There is a clear need for better testing options and education for clinicians who treat
49
50 suspected synthetic cathinone overdose. ED clinicians must rely heavily on self-reports to
51
52 determine the cause of a drug overdose, in part because urine toxicology tests can take days to
53
54 complete. Vazirian et al. (2015) surveyed ED clinicians and found that 77% do not explicitly ask
55
56 if a patient has taken synthetic cathinones, and 98% were unaware of bath salt urine toxicology
57
58
59
60
61
62
63
64
65

1
2
3
4 tests. While 60% did report having encountered a patient who admitted to bath salts use, these
5
6 patients presented with symptoms that could have easily been misattributed to more common
7
8 drugs of abuse. These symptoms include agitation (81%), aggression or violence (65%), or
9
10 hallucinations (46%) – all common effects of the widely used drugs that synthetic cathinones
11
12 may be intended to mimic.
13
14

15 16 **Unintentional Synthetic Cathinone Use, Combinatorial Effects, and Underlying** 17 18 **Neurobiology.** 19

20
21 Unintentional synthetic cathinone use greatly increases the risk of overdose and death.
22
23 The two most common causes of death from synthetic cathinone use are overdose and fatal self-
24
25 harm (citation below). Our speculation is that overdoses occur because of a fundamental
26
27 misunderstanding of the mechanisms and potencies of the synthetic cathinone(s) being used.
28
29 Self-harm associated with synthetic cathinone use may result from serious psychological effects
30
31 of intoxication such as loss of impulse control, increased risk-taking/sensation-seeking,
32
33 hallucinations, and paranoia (_____). Moreover, chronic use can lead to persistent
34
35 neurochemical alterations that underlie extreme depression, thereby triggering self-harm (Banks
36
37 et al. 2014; Barrio et al. 2016).
38
39
40
41

42
43 In general, the effects of synthetic cathinones are attributed to the neurotransmitters that
44
45 most illicit drugs act upon including dopamine (DA), noradrenaline (NA) and serotonin (5-HT).
46
47 Drugs such as cocaine function as DA and NA transporter (reuptake) blockers leading to net
48
49 elevations in synaptic neurotransmitter levels. Compared to cocaine, MDPV increases synaptic
50
51 NA with approximately 10-fold- and synaptic DA with approximately 50-fold-greater potency
52
53 (Baumann et al. 2012). Therefore, if one consumes MDPV under the impression that it is
54
55 cocaine, the resulting intoxication would be manifold greater than intended. Excitotoxic levels of
56
57
58
59
60
61
62
63
64
65

1
2
3
4 DA and NA produce hyperthermia and increased heart rate that can lead to multiorgan failure
5
6 and death (Walter and Carretto 2015).
7

8
9 Drugs such as MDMA, amphetamine, and methamphetamine act as DA, 5-HT, and NA
10 transporter substrates, meaning the drug enters the cell and disrupts vesicular storage and
11
12 transmitter release dynamics (e.g., Baumann et al. 2013). Transporter substrates produce
13
14 persistent depletion of neurotransmitters and loss of functional transporters (Baumann et al.
15
16 2007, 2013; Gygi et al. 1997; German et al. 2014). Indeed depression and suicide are particularly
17
18 common in heavy users of both MDMA and the mechanistically similar synthetic cathinone,
19
20 mephedrone (Cohen 1996; Marinetti and Antonides 2013; Elliott and Evans 2014).
21
22
23
24
25

26 Death from unintentional synthetic cathinone overdose is typified by one of the first
27 reported cases following unintentional “bath salts” consumption. Warrick et al. (2012) describe a
28
29 24-year-old female who died after consuming two pills of what turned out to be high doses of
30
31 methylone and butylone rather than the “ecstasy” she believed she was taking. The patient was
32
33 found unconscious at a concert and was febrile, tachypnic, and hypertensive. An ED exam
34
35 revealed that she was diaphoretic, tremulous, hyperflexic, and had sustained clonus. Despite
36
37 maximal supportive care, optimal ventilation, and improvements in hemodynamics and
38
39 coagulopathy, within 48 hours of admission the patient developed acute respiratory distress
40
41 syndrome, renal failure, hypoxemia, lactic acidosis and ultimately expired. Urinalysis and
42
43 GC/MS testing of pills found on her person similar to the two ingested revealed the sole presence
44
45 of methylone and butylone, with no traces of MDMA that the “ecstasy” was sold as. While
46
47 methylone and butylone are 2-6x less potent than MDMA in terms of DA and 5-HT transporter
48
49 substrate and blocking properties, the excessive quantity of each compound within each pill are
50
51 the likely reason the patient expired. Ecstasy tablets typically contain 30-150 mg of MDMA, and
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 while methylone and butylone are up to 10x less effective at reducing 5-HT and NA, the three
5
6 drugs are similarly effective at increasing DA which may explain why the three drugs are
7
8 consumed recreationally at similar doses (e.g., Hall and Henry 2006; López-Arnau et al. 2012;
9
10 Warrick et al. 2012). In the case described by Warrick et al. (2012), each pill contained 422 mg
11
12 of methylone and 53 mg of butylone - doses that would produce profoundly higher DA levels
13
14 than what is considered pleasurable or safe.
15
16
17

18
19 Unintentional synthetic cathinone use can occur in polydrug users, and a small number of
20
21 preclinical studies have begun to elucidate the neurotoxic and behavioral effects of drug
22
23 combinations. Table 1 lists effects observed in animal subjects following exposure to drug
24
25 combinations that include a synthetic cathinone. Notably, enhancement of neurotoxicity is
26
27 observed with drugs that have overlapping substrate properties (e.g., mephedrone elevates
28
29 MDMA-induced DA neurotoxicity; Angoa-Perez 2013). Some combinations, however, produce
30
31 unexpected protective effects that may be due to the mechanism of one drug minimizing effects
32
33 of a second drug. For example, the DA transporter blocking properties of MDPV may protect
34
35 against the DA releasing properties of methamphetamine (Anneken et al. 2015). In addition,
36
37 many studies fail to account for potential contributions of temperature when examining the
38
39 neurotoxic effects of synthetic cathinone use (alone or in combination with a second drug). For
40
41 example, mephedrone potentiates the neurotoxic effects of MDMA under group- but not single-
42
43 housing conditions (Hadlock et al. 2011) which likely reflects the inherent risk of consuming
44
45 these combinations in crowded environments (e.g. EDM clubs). The neurotoxicity observed by
46
47 Hadlock et al. (2011) was persistent yet limited to cells producing 5-HT. In humans, 5-HT
48
49 depletion has been associated with the negative mood, irritability, and suicidal thoughts
50
51 experienced in the days after MDMA or mephedrone use (Cohen 1996; Freeman et al. 2012).
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 More research will be needed to reach a clearer understanding of how synthetic cathinones
5
6 impact the neurochemical and behavioral effects experienced from other drugs when provided,
7
8 knowingly or unbeknownst to the consumer, in polydrug formulations.
9

10
11 Synthetic cathinone users are at additional risk of experiencing psychosis. There are
12
13 several reported cases of years-long, intermittent mephedrone users entering drug rehabilitation
14
15 for delusional thoughts and auditory and visual hallucinations (Barrio 2016). Psychosis can occur
16
17 in patients with no medical or family history of mental illness demonstrating the profound impact
18
19 of long-term drug use that may have been consumed unknowingly. MDPV can produce delirium
20
21 - a state characterized by bizarre behavior, paranoia, and hallucinations. For example, Penders
22
23 and Gestring (2011) describe 3 cases of individuals experiencing delusions of persecution and
24
25 harassment from nonexistent sources: one man hallucinated his estranged wife tapping on his
26
27 window, a young housewife was filmed describing home invaders in an unoccupied room, and a
28
29 middle-aged man hallucinated burglars shooting him with laser beams. Moreover, the memory
30
31 disturbances common in MDPV-induced delirium greatly increase the risk of overdose death
32
33 because patients frequently have no recollection of the amount, frequency, or route of
34
35 administration. This is especially dangerous in the case of unintentional use of MDPV, of a drug
36
37 that is over ten times more potent than the more common substance it may have been intended to
38
39 mimic (e.g. cocaine). These cases demonstrate the great risk involved in consuming a substance
40
41 that may be far more potent than one conceives.
42
43
44
45
46
47
48
49

50 Finally, unintentional synthetic cathinone use carries with it significant risk of death from
51
52 self-harm. Self-harm has been cited as the second most common cause of death in synthetic
53
54 cathinone users (Schifano et al. 2012; Marinetti and Antonides 2013). A 3-year review of novel
55
56 psychoactive substances in casework (Elliott and Evans 2014) found that in deaths following
57
58
59
60
61
62
63
64
65

1
2
3
4 synthetic cathinone use, 41% were hangings or other mechanical suicide (i.e., fatal self-harm),
5
6 which was the highest proportion compared to other classes of drugs. While hangings are the
7
8 most common form of fatal self-harm, drownings, gunshots wounds, repeated self-lacerations,
9
10 and jumping from bridges have all been reported (Schifano et al. 2012; Marinetti and Antonides
11
12 2013).
13
14

15
16 Self-harm related to synthetic cathinone use is attributed to impulsivity, paranoia, and
17
18 violent behavior in response to vivid hallucinations or delusional thoughts (Penders 2012;
19
20 Penders and Gestring 2011). In a case of fatal self-harm associated with synthetic cathinone use,
21
22 a 21-year old with no history of depression committed suicide 5 days after ingesting an unknown
23
24 powder later determined to be MDPV (PBS News, 2018). The decedent experienced days of
25
26 insomnia and a frightening hallucination of 25 police cars outside his kitchen. Upon being told
27
28 the police cars did not exist, the patient slit his throat in front of his parents. The self-inflicted
29
30 laceration was non-fatal and was successfully stitched closed at a hospital. However, 4 hours
31
32 after returning home from the hospital, and 5 days after initial drug-taking, the decedent
33
34 committed suicide with a rifle. This case exemplifies how MDPV use can promote self-harm and
35
36 suicide, possibly as a consequence of withdrawal-associated affective distress. It is likely the
37
38 decedent was experiencing excitotoxic levels of DA during the psychotic episode. Synthetic
39
40 cathinone-induced DA excitotoxicity can take days to normalize (Den Hollander et al. 2013;
41
42 Martínez-Clemente et al. 2014), during which depression and suicidal thoughts may emerge.
43
44 Indeed, the term “suicide tuesday” emerged to describe the depressive withdrawal state
45
46 experienced days following initial drug-taking (Psychedelics.com 2018). It is possible that
47
48 antipsychotic treatment may have prevented death of the patient, as antipsychotic drugs working
49
50 as DA modulators synthetic cathinone-induced psychosis (Banks et al. 2014; Barrio et al. 2016).
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 ED clinicians and potential drug users can reduce the risk of overdose and death from
5
6 unintentional synthetic cathinone use. Clinicians should be aware of the constellation of
7
8 symptoms typical of synthetic cathinone overdose: agitation, sustained hyperthermia, muscle
9
10 rigidity, and psychosis. The typical course of treatment involves aggressive cooling, hydration,
11
12 and antipsychotic medication administration. Additionally, drug users who reject abstinence can
13
14 make use of pill-testing services that can reliably detect the presence of ‘unknown substances’ in
15
16 drugs such as cocaine and ecstasy (Saleemi et al. 2017). Drug users and clinicians should also
17
18 note the potential for self-harm and suicide in patients during intoxication and during
19
20 withdrawal. Future studies examining the toxicity associated with synthetic cathinone drug
21
22 combinations are also needed to better address the emerging phenomenon of unintentional
23
24 synthetic cathinone use.
25
26
27
28
29

30 31 **Concluding Remarks.** 32

33 Adulteration of illegal drugs with synthetic cathinones is an untenable yet largely
34
35 unrecognized public health concern, the magnitude of which we are only beginning to appreciate
36
37 through empirical observation. Traditional approaches used to assess the effects of drugs on
38
39 public health and safety (e.g., surveys, urinalysis) are likely underestimating synthetic cathinone
40
41 use, especially in high-risk populations. Moreover, synthetic cathinone abuse carries with it risks
42
43 unique to this class of drugs such as overdose, self-harm, and suicide. It is clear that synthetic
44
45 cathinones mimicking relatively popular illegal drugs such as cocaine (e.g., MDPV) and MDMA
46
47 (e.g., methyldone and mephedrone) produce similar euphorogenic subjective effects, though
48
49 typically at fractions of the dose.
50
51
52
53
54

55 We believe the burden of addressing this public health problem is complicated and shared
56
57 by many. In effort to specifically reduce accidental overdose, we advocate the development and
58
59
60
61
62
63
64
65

1
2
3
4 commercial sale of purity detectors (e.g. pill-testing products and services) that can be employed
5
6 by users. We advocate the dissemination of accurate, empirically-derived information on the
7
8 effects of synthetic cathinones so users and medical professionals including ED personnel can
9
10 recognize when an individual - knowingly or not - has consumed a synthetic cathinone.
11
12
13 Moreover, ED personnel should be aware of the availability of synthetic cathinone drug-testing
14
15 products and services. In the future, collaborative work between field observers and basic
16
17 scientists will (i) clarify the extent to which synthetic cathinones impact the subjective and
18
19 physiological effects associated with other drug use, (ii) determine how situational factors such
20
21 as environment influence the effects of synthetic cathinones taken with other drugs, and (iii) aide
22
23 in the development of medical intervention strategies associated with the unique risks of drug use
24
25 complicated by synthetic cathinone adulteration.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57

58 **References.**

59
60
61
62
63
64
65

Alles GA, Fairchild MD, Jensen M (1961) Chemical pharmacology of Catha edulis. J Med Pharm Chem 3:323–52

Anneken JH, Angoa-Pérez M, Kuhn DM (2015) 3,4-Methylenedioxypyrovalerone prevents while methylone enhances methamphetamine-induced damage to dopamine nerve endings: β -ketoamphetamine modulation of neurotoxicity by the dopamine transporter. J Neurochem 133:211–222. doi: 10.1111/jnc.13048

Alvarez J-C, Fabresse N, Larabi IA (2018) Prevalence and Surveillance of Synthetic Cathinones Use by Hair Analysis: An Update Review. Curr Pharm Des 23:5487–5495. doi: 10.2174/1381612823666170704124156

American Association of Poison Control Centers 2016. Bath salts data.

https://aapcc.s3.amazonaws.com/files/library/Bath_Salts_Web_Data_through_3.2016.pdf

. Accessed on 30 November 2018

Banks ML, Worst TJ, Rusyniak DE, Sprague JE (2014) Synthetic Cathinones (“Bath Salts”). J Emerg Med 46:632–642. doi: 10.1016/j.jemermed.2013.11.104

Barratt MJ, Bruno R, Ezard N, Ritter A (2018) Pill testing or drug checking in Australia: Acceptability of service design features. Drug Alcohol Rev 37:226–236. doi: 10.1111/dar.12576

Barrio P, Gaskell M, Goti J, et al (2016) Persistent psychotic symptoms after long-term heavy use of mephedrone: A two-case series. Adicciones 28:154–7

Baumann MH, Partilla JS, Lehner KR, et al (2012) Powerful Cocaine-Like Actions of 3,4-Methylenedioxypyrovalerone (MDPV), a Principal Constituent of Psychoactive ‘Bath Salts’ Products TL - 38. Neuropsychopharmacology 38 VN-r:552–562. doi: 10.1038/npp.2012.204

- 1
2
3
4 Baumann MH, Partilla JS, Lehner KR, et al (2013) Powerful Cocaine-Like Actions of 3,4-
5
6 Methylenedioxypyrovalerone (MDPV), a Principal Constituent of Psychoactive ‘Bath
7
8 Salts’ Products. *Neuropsychopharmacology* 38:552–562. doi: 10.1038/npp.2012.204
9
10
11 Brunt TM, Atkinson AM, Nefau T, et al (2017) Online test purchased new psychoactive
12
13 substances in 5 different European countries: A snapshot study of chemical composition
14
15 and price. *Int J Drug Policy* 44:105–114. doi: 10.1016/j.drugpo.2017.03.006
16
17
18 Brunt TM, Poortman A, Niesink RJ, van den Brink W (2011a) Instability of the ecstasy market
19
20 and a new kid on the block: mephedrone. *J Psychopharmacol* 25:1543–1547. doi:
21
22 10.1177/0269881110378370
23
24
25 Brunt TM, Poortman A, Niesink RJ, van den Brink W (2011b) Instability of the ecstasy market
26
27 and a new kid on the block: mephedrone. *J Psychopharmacol* 25:1543–1547. doi:
28
29 10.1177/0269881110378370
30
31
32
33 Butterfield RJ, Barratt MJ, Ezard N, Day RO (2016) Drug checking to improve monitoring of
34
35 new psychoactive substances in Australia. *Med J Aust* 204:144–5
36
37
38 Carbone PN, Carbone DL, Carstairs SD, Luzi SA (2013) Sudden cardiac death associated with
39
40 methylone use. *Am J Forensic Med Pathol* 34:26–8. doi:
41
42 10.1097/PAF.0b013e31827ab5da
43
44
45 Caudevilla-Gálligo F, Ventura M, Indave Ruiz BI, Fornís I (2013) Presence and composition of
46
47 cathinone derivatives in drug samples taken from a Drug Test Service in Spain (2010-
48
49 2012). *Hum Psychopharmacol Clin Exp* 28:341–344. doi: 10.1002/hup.2296
50
51
52
53 CBS (2016) After ravaging Florida, street drug flakka disappears - CBS News. In: CBS News.
54
55 <https://www.cbsnews.com/news/after-ravaging-florida-street-drug-flakka-disappears/>.
56
57
58 Accessed 28 May 2018
59
60
61
62
63
64
65

- 1
2
3
4 Cohen RS (1996) Adverse symptomatology and suicide associated with the use of
5
6 methylenedioxymethamphetamine (MDMA; “Ecstasy”). *Biol Psychiatry* 39:819–820.
7
8 doi: 10.1016/0006-3223(95)00551-X
9
- 10
11 Den Hollander B, Rozov S, Linden A-M, et al (2013) Long-term cognitive and neurochemical
12
13 effects of “bath salt”; designer drugs methylone and mephedrone. doi:
14
15 10.1016/j.pbb.2012.10.006
16
17
- 18
19 Dorairaj JJ, Healy C, McMenamin M, Eadie PA (2012) The untold truth about “bath salt”; highs:
20
21 A case series demonstrating local tissue injury. *J Plast Reconstr Aesthet Surg* 65:e37-41.
22
23 doi: 10.1016/j.bjps.2011.10.004
24
25
- 26
27 Drug Enforcement Administration U, Control Division D (2015) National Forensic Laboratory
28
29 Information System: 2016 Annual Report
30
- 31
32 Drug Enforcement Agency (2017) Drug of Abuse: A DEA Resource Guide 2017
33
34 Edition
35
- 36
37 Elliott S, Evans J (2014) A 3-year review of new psychoactive substances in casework. *Forensic*
38
39 *Sci Int* 243:55–60. doi: 10.1016/J.FORSCIINT.2014.04.017
40
- 41
42 European Monitoring Centre for Drugs and Drug Addiction (2015) European Drug Report 2015.
43
44 doi: 10.2810/084165
45
- 46
47 Fernández-Calderón F, Cleland CM, Palamar JJ (2018) Polysubstance use profiles among
48
49 electronic dance music party attendees in New York City and their relation to use of new
50
51 psychoactive substances. *Addict Behav* 78:85–93. doi: 10.1016/j.addbeh.2017.11.004
52
- 53
54 Freeman TP, Morgan CJA, Vaughn-Jones J, et al (2012) Cognitive and subjective effects of
55
56 mephedrone and factors influencing use of a ‘new legal high.’ *Addiction* 107:792–800.
57
58 doi: 10.1111/j.1360-0443.2011.03719.x
59
60
61
62
63
64
65

German CL, Fleckenstein AE, Hanson GR (2014) Bath salts and synthetic cathinones: an emerging designer drug phenomenon. *Life Sci* 97:2–8. doi: 10.1016/j.lfs.2013.07.023

Glennon RA, Yousif M, Naiman N, Kalix P (1987) Methcathinone: a new and potent amphetamine-like agent. *Pharmacol Biochem Behav* 26:547–51

Gygi MP, Fleckenstein AE, Gibb JW, Hanson GR (1997) Role of endogenous dopamine in the neurochemical deficits induced by methcathinone. *J Pharmacol Exp Ther* 283:1350–5

Hyde JF, Browning E, Adams R (1928) Synthetic Homologs of d,l-ephedrine. *J Am Chem Soc* 50:2287–2292. doi: 10.1021/ja01395a032

Johnston LD, Miech RA, O'malley PM, et al (2018) Key Findings on Adolescent Drug Use

Kalix P (1984) The pharmacology of khat. *Gen Pharmacol Vasc Syst* 15:179–187. doi: 10.1016/0306-3623(84)90156-3

Kelly BC, Wells BE, Pawson M, et al (2013) Novel psychoactive drug use among younger adults involved in US nightlife scenes. *Drug Alcohol Rev* 32:588–593. doi: 10.1111/dar.12058

Kintz P, Salomone A, Vincenti M (2015) Hair analysis in clinical and forensic toxicology

Krotulski AJ, Papsun DM, De Martinis BS, et al (2018) N-Ethyl Pentylone (Ephylone) Intoxications: Quantitative Confirmation and Metabolite Identification in Authentic Human Biological Specimens. *J Anal Toxicol*. doi: 10.1093/jat/bky025

Kurtz SP, Buttram ME, Pagano ME, Surratt HL (2017) A randomized trial of brief assessment interventions for young adults who use drugs in the club scene. *J Subst Abuse Treat* 78:64–73. doi: 10.1016/j.jsat.2017.05.008

Majchrzak M, Celiński R, Kuś P, et al (2018) The newest cathinone derivatives as designer drugs: an analytical and toxicological review. *Forensic Toxicol* 36:33–50. doi: 10.1007/s11419-017-0385-6

1
2
3
4 Marchei E, Pacifici R, Mannocchi G, et al (2018) New synthetic opioids in biological and non-
5
6 biological matrices: A review of current analytical methods. *TrAC Trends Anal Chem*
7
8 102:1–15. doi: 10.1016/J.TRAC.2018.01.007
9

10
11 Marinetti LJ, Antonides HM (2013) Analysis of synthetic cathinones commonly found in bath
12
13 salts in human performance and postmortem toxicology: method development, drug
14
15 distribution and interpretation of results. *J Anal Toxicol* 37:135–46. doi:
16
17 10.1093/jat/bks136
18
19

20
21 Martínez-Clemente J, López-Arnau R, Abad S, et al (2014) Dose and time-dependent selective
22
23 neurotoxicity induced by mephedrone in mice. *PLoS One* 9:e99002. doi:
24
25 10.1371/journal.pone.0099002
26
27

28
29 Martin M, Muller JF, Turner K, et al (2012) Evidence of mephedrone chronic abuse through hair
30
31 analysis using GC/MS. *Forensic Sci Int* 218:44–8. doi: 10.1016/j.forsciint.2011.10.016
32
33

34
35 Miller BL, Stogner JM (2014) Not-So-Clean Fun: A Profile of Bath Salt Users Among a College
36
37 Sample in the United States. *J Psychoactive Drugs* 46:147–153. doi:
38
39 10.1080/02791072.2013.876520
40

41
42 National Forensic Information System (2011) New Methodology for Calculating National and
43
44 Regional Estimates and Presenting Data in Publications
45

46
47 Newshubb Staff (2018) 13 hospitalised in Christchurch after taking drugs they thought were
48
49 MDMA | Newshub. In: Newshub. [http://www.newshub.co.nz/home/new-](http://www.newshub.co.nz/home/new-zealand/2018/03/13-hospitalised-in-christchurch-after-taking-drugs-they-thought-were-mdma.html)
50
51 [zealand/2018/03/13-hospitalised-in-christchurch-after-taking-drugs-they-thought-were-](http://www.newshub.co.nz/home/new-zealand/2018/03/13-hospitalised-in-christchurch-after-taking-drugs-they-thought-were-mdma.html)
52
53 [mdma.html](http://www.newshub.co.nz/home/new-zealand/2018/03/13-hospitalised-in-christchurch-after-taking-drugs-they-thought-were-mdma.html). Accessed 24 May 2018
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 NIDA (2018) Synthetic Cathinones (" Bath Salts "). In: NIDA Drug Facts.

5
6
7 <https://www.drugabuse.gov/publications/drugfacts/synthetic-cathinones-bath-salts>.

8
9 Accessed 26 May 2018

10
11 Nóbrega L, Dinis-Oliveira RJ (2018) The synthetic cathinone α -pyrrolidinovalerophenone (α -

12
13 PVP): pharmacokinetic and pharmacodynamic clinical and forensic aspects. *Drug Metab*

14
15
16 *Rev* 50:125–139. doi: 10.1080/03602532.2018.1448867

17
18
19 Numan N (2004) Exploration of adverse psychological symptoms in Yemeni khat users by the

20
21 *Symptoms Checklist-90 (SCL-90)*. *Addiction* 99:61–5

22
23
24 Palamar JJ (2017) There's something about Molly: The underresearched yet popular powder

25
26 form of ecstasy in the United States. *Subst Abus* 38:15–17. doi:

27
28 10.1080/08897077.2016.1267070

29
30
31 Palamar JJ (2018) "Bath Salt" Use and Beliefs about Use among Electronic Dance Music

32
33 Attendees. *J Psychoactive Drugs* 1–8. doi: 10.1080/02791072.2018.1517229

34
35
36 Palamar JJ (2015) "Bath Salt"; use among a nationally representative sample of high school

37
38 seniors in the United States. *Am J Addict* 24:488–91. doi: 10.1111/ajad.12254

39
40
41 Palamar JJ, Acosta P, Calderón FF, et al (2017a) Assessing self-reported use of new

42
43 psychoactive substances: The impact of gate questions. *Am J Drug Alcohol Abuse*

44
45 43:609–617. doi: 10.1080/00952990.2017.1322094

46
47
48 Palamar JJ, Acosta P, Cleland CM (2018) Attitudes and Beliefs About New Psychoactive

49
50 Substance Use Among Electronic Dance Music Party Attendees. *Subst Use Misuse*

51
52 53:381–390. doi: 10.1080/10826084.2017.1327980

53
54
55

56
57
58

59
60
61

62
63
64

65

- 1
2
3
4 Palamar JJ, Acosta P, Sherman S, et al (2016a) Self-reported use of novel psychoactive
5
6 substances among attendees of electronic dance music venues. *Am J Drug Alcohol Abuse*
7
8 42:624–632. doi: 10.1080/00952990.2016.1181179
9
10
11 Palamar JJ, Barratt MJ, Ferris JA, Winstock AR (2016b) Correlates of new psychoactive
12
13 substance use among a self-selected sample of nightclub attendees in the United States.
14
15 *Am J Addict* 25:400–407. doi: 10.1111/ajad.12403
16
17
18 Palamar JJ, Griffin-Tomas M, Ompad DC (2015a) Illicit drug use among rave attendees in a
19
20 nationally representative sample of US high school seniors. *Drug Alcohol Depend*
21
22 152:24–31. doi: 10.1016/j.drugalcdep.2015.05.002
23
24
25
26 Palamar JJ, Martins SS, Su MK, Ompad DC (2015b) Self-reported use of novel psychoactive
27
28 substances in a US nationally representative survey: Prevalence, correlates, and a call for
29
30 new survey methods to prevent underreporting. *Drug Alcohol Depend* 156:112–119. doi:
31
32 10.1016/j.drugalcdep.2015.08.028
33
34
35
36 Palamar JJ, Salomone A, Gerace E, et al (2017b) Hair testing to assess both known and unknown
37
38 use of drugs amongst ecstasy users in the electronic dance music scene. *Int J Drug Policy*
39
40 48:91–98. doi: 10.1016/j.drugpo.2017.07.010
41
42
43 Palamar JJ, Salomone A, Vincenti M, Cleland CM (2016c) Detection of “bath salts”; and other
44
45 novel psychoactive substances in hair samples of ecstasy/MDMA/”Molly”; users. *Drug*
46
47 *Alcohol Depend* 161:200–5. doi: 10.1016/j.drugalcdep.2016.02.001
48
49
50 Palamar JJ, Salomone A, Vincenti M, Cleland CM (2016d) Detection of “bath salts”; and other
51
52 novel psychoactive substances in hair samples of ecstasy/MDMA/”Molly”; users. *Drug*
53
54 *Alcohol Depend* 161:200–5. doi: 10.1016/j.drugalcdep.2016.02.001
55
56
57
58
59
60
61
62
63
64
65

Papaseit E, Pérez-Mañá C, Mateus J-A, et al (2016) Human Pharmacology of Mephedrone in Comparison with MDMA. *Neuropsychopharmacology* 41:2704–2713. doi: 10.1038/npp.2016.75

PBS News The Drug That Never Lets Go | PBS NewsHour | Sept. 20, 2012 | PBS.
<https://www.pbs.org/newshour/spc/multimedia/bath-salts/>. Accessed 28 Sep 2018

Penders TM (2012) How to recognize a patient who's high on "bath salts". *J Fam Pract* 61:210–2

Penders TM, Gestring R (2011) Hallucinatory delirium following use of MDPV: "Bath Salts." *Gen Hosp Psychiatry* 33:525–526. doi: 10.1016/j.genhosppsych.2011.05.014

Psychedelics.com What Is Suicide Tuesday? <http://psychedelics.com/mdma/what-is-suicide-tuesday/>. Accessed 28 Sep 2018

Ridpath A, Driver CR, Nolan ML, et al (2014) Illnesses and deaths among persons attending an electronic dance-music festival - New York City, 2013. *MMWR Morb Mortal Wkly Rep* 63:1195–1198

Ross MW, Mattison AM, Franklin DR (2003) Club drugs and sex on drugs are associated with different motivations for gay circuit party attendance in men. *Subst Use Misuse* 38:1173–83

Rust KY, Baumgartner MR, Dally AM, Kraemer T (2012) Prevalence of new psychoactive substances: A retrospective study in hair. *Drug Test Anal* 4:402–408. doi: 10.1002/dta.1338

Saleemi S, Pennybaker SJ, Wooldridge M, Johnson MW (2017) Who is 'Molly'? MDMA adulterants by product name and the impact of harm-reduction services at raves. *J Psychopharmacol* 31:1056–1060. doi: 10.1177/0269881117715596

1
2
3
4 Salomone A, Gazzilli G, Di Corcia D, et al (2016a) Determination of cathinones and other
5
6 stimulant, psychedelic, and dissociative designer drugs in real hair samples. Anal Bioanal
7
8 Chem 408:2035–2042. doi: 10.1007/s00216-015-9247-4
9
10
11 Salomone A, Gazzilli G, Di Corcia D, et al (2016b) Determination of cathinones and other
12
13 stimulant, psychedelic, and dissociative designer drugs in real hair samples. Anal Bioanal
14
15 Chem 408:2035–2042. doi: 10.1007/s00216-015-9247-4
16
17
18 Salomone A, Palamar JJ, Gerace E, et al (2017) Hair Testing for Drugs of Abuse and New
19
20 Psychoactive Substances in a High-Risk Population. J Anal Toxicol 41:376–381. doi:
21
22 10.1093/jat/bkx020
23
24
25 Schifano F, Corkery J, Ghodse AH (2012) Suspected and Confirmed Fatalities Associated With
26
27 Mephedrone (4-Methylmethcathinone, “Meow Meow”) in the United Kingdom. J Clin
28
29 Psychopharmacol 32:710–714. doi: 10.1097/JCP.0b013e318266c70c
30
31
32 State Department (1994) Federal Register: Document number 94-14962.
33
34 <https://www.gpo.gov/fdsys/pkg/FR-1994-12-01/html/94-29562.htm>. Accessed 24 May
35
36 2018
37
38
39 Substance Abuse and Mental Health Services Administration S (2011) The DAWN Report Data
40
41 Spotlight: “Bath Salts” Were Involved in over 20,000 Drug-Related Emergency
42
43 Department Visits in 2011
44
45
46 Vazirian M, Jerry JM, James J, Dale RM (2015) Bath Salts in the Emergency Department. J
47
48 Addict Med 9:94–98. doi: 10.1097/ADM.0000000000000094
49
50
51 Walter E, Carraretto M (2015) Drug-induced hyperthermia in critical care. J Intensive Care Soc
52
53 16:306–311. doi: 10.1177/1751143715583502
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 Warrick BJ, Wilson J, Hedge M, et al (2012) Lethal Serotonin Syndrome After Methylone and
5
6 Butylone Ingestion. J Med Toxicol 8:65–68. doi: 10.1007/s13181-011-0199-6
7
8

9 Wood DM, Davies S, Puchnarewicz M, et al (2010) Recreational Use of Mephedrone (4-
10
11 Methylmethcathinone, 4-MMC) with Associated Sympathomimetic Toxicity. J Med
12
13 Toxicol 6:327–330. doi: 10.1007/s13181-010-0018-5
14
15

16 Wood KE (2013) Exposure to Bath Salts and Synthetic Tetrahydrocannabinol from 2009 to 2012
17
18 in the United States. J Pediatr 163:213–216. doi: 10.1016/j.jpeds.2012.12.056
19
20


21 Zuba D, Adamowicz P (2018) Analytical Methods Used for Identification and Determination of
22
23 Synthetic Cathinones and Their Metabolites. Springer, Cham, pp 41–69
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Tables.

Table 1. Neurotoxic and Behavioral Effects of Synthetic Cathinones in Combination More Commonly Used Illegal Drugs. Synthetic cathinones can potentiate, nullify, or protect against the neurotoxic and behavioral effects of more commonly used drugs (second drug).

Synthetic cathinone	Common Illegal Drug	Effects (neurotoxic or behavioral)	Reference (format)
Mephedrone	MDMA	Potentiates MDMA-induced DA and 5-HT neurotoxicity in group-housed animals	Hadlock et al. 2011
MDPV	Methamphetamine	Reduced methamphetamine-induced DA neurotoxicity	Anneken 2015
Methylone			
Methylone	MDMA	Reduced MDMA-elicited astrogliosis and neurotoxicity markers	Miner 2017
Mephedrone	MDMA	No effect on 5-HT neurotoxicity	Angoa-Perez 2014
	Methamphetamine		
Mephedrone	MDMA	Reduced depression, improved memory consolidation	Budzynska 2017
Mephedrone	Methamphetamine	Enhanced DA neurotoxicity	Angoa-Perez 2013
	Amphetamine		
	MDMA		



[Click here to access/download](#)

Supplementary Material

Signed Scanned Oliver COI form.pdf

